

S_4 Flavor Symmetry of Quarks and Leptons in SU(5) GUT

Hajime Ishimori^{1,*}, Yusuke Shimizu^{1,†},
Morimitsu Tanimoto^{2,‡}

¹*Graduate School of Science and Technology, Niigata University,
Niigata 950-2181, Japan*

²*Department of Physics, Niigata University, Niigata 950-2181, Japan*

Abstract

We present a S_4 flavor model to unify quarks and leptons in the framework of the SU(5) GUT. Three generations of $\bar{5}$ -plets in SU(5) are assigned 3_1 of S_4 while the first and the second generations of 10-plets in SU(5) are assigned to be 2 of S_4 , and the third generation of 10-plet is to be 1_1 of S_4 . Right-handed neutrinos are also assigned 2 for the first and second generations and 1_1 for the third generation, respectively. Taking vacuum alignments of relevant gauge singlet scalars, we predict the quark mixing as well as the tri-bimaximal mixing of neutrino flavors. Especially, the Cabibbo angle is predicted to be 15° in the limit of the vacuum alignment. We can improve the model to predict observed CKM mixing angles as well as the non-vanishing U_{e3} of the neutrino flavor mixing.

*E-mail address: ishimori@muse.sc.niigata-u.ac.jp

†E-mail address: shimizu@muse.sc.niigata-u.ac.jp

‡E-mail address: tanimoto@muse.sc.niigata-u.ac.jp

1 Introduction

Neutrino experimental data provide an important clue for elucidating the origin of the observed hierarchies in mass matrices for quarks and leptons. Recent experiments of the neutrino oscillation go into a new phase of precise determination of mixing angles and mass squared differences [1], which indicate the tri-bimaximal mixing for three flavors in the lepton sector [2]. These large mixing angles are completely different from the quark mixing ones. Therefore, it is very important to find a natural model that leads to these mixing patterns of quarks and leptons with good accuracy.

Flavor symmetry is expected to explain the mass spectrum and the mixing matrix of quarks and leptons. In particular, some predictive models with non-Abelian discrete flavor symmetries have been explored by many authors. Among them, the tri-bimaximal mixing of leptons has been understood based on the non-Abelian finite group S_3 [3]-[19], A_4 [20]-[41], and T' [42]-[47], because these symmetries provide the definite meaning of generations and connects different generations. On the other hand, much attention has been devoted to the question whether these models can be extended to describe the observed pattern of quark masses and mixing angles, and whether these can be made compatible with the SU(5) or SO(10) grand unified theory (GUT).

Recently, group-theoretical arguments indicate that the discrete symmetry S_4 is the minimal flavor symmetry compatible with the tri-bimaximal neutrino mixing [48]. Actually, the exact tri-bimaximal neutrino mixing is realized in the S_4 flavor model [49]. Thus, the S_4 flavor model is attractive for the lepton sector [50]-[54]. Although an attempt to unify the quark and lepton sector was presented towards a grand unified theory of flavor [51], mixing angles are not predicted clearly.

In our work, we present a S_4 flavor model to unify the quarks and leptons in the framework of the SU(5) GUT. The group S_4 has irreducible representations 1_1 , 1_2 , 2 , 3_1 , and 3_2 . Three generations of $\bar{5}$ -plets in SU(5) are assigned 3_1 of S_4 while the first and the second generations of 10-plets in SU(5) are assigned to be 2 of S_4 , and the third generation of 10-plet is to be 1_1 of S_4 . These assignments of S_4 for $\bar{5}$ and 10 lead to the completely different structure of quark and lepton mass matrices. Right-handed neutrinos, which are SU(5) gauge singlets, are also assigned 2 for the first and second generations, and 1_1 for the third generation, respectively. These assignments are essential to realize the tri-bimaximal mixing of neutrino flavors. Taking vacuum alignments of relevant gauge singlet scalars, we predict the quark mixing as well as the tri-bimaximal mixing of leptons. Especially, the Cabibbo angle is predicted to be 15° in the limit of the vacuum alignment. We improve the model to predict observed CKM mixing angles as well as the non-vanishing U_{e3} of the neutrino flavor mixing.

The paper is organized as follows. We present the prototype of the S_4 flavor model of quarks and leptons in SU(5) GUT in section 2, and discuss the lepton sector in section 3, and the quark sector in section 4. In section 5, we present the improved model with additional scalar in order to study detail of the model. Section 6 is devoted to the summary. In the appendix, we present the multiplication rules of S_4 , and the scalar potential analysis. Vacuum alignments and magnitude of VEVs are also summarized in the appendix.

2 Prototype of S_4 flavor model in SU(5) GUT

We present the prototype of the S_4 flavor model in the SU(5) GUT to understand the essence of our model clearly. We consider the supersymmetric GUT based on SU(5). The flavor symmetry of quarks and leptons is the discrete group S_4 in our model. The group S_4 has irreducible representations 1_1 , 1_2 , 2 , 3_1 , and 3_2 . The multiplication rules of S_4 are summarized in appendix.

	T_3	(T_1, T_2)	(F_1, F_2, F_3)	(N_e^c, N_μ^c)	N_τ^c	H_5	$H_{\bar{5}}$
SU(5)	10	10	$\bar{5}$	1	1	5	$\bar{5}$
S_4	1_1	2	3_1	2	1_1	1_1	1_1
Z_4	ω^2	ω^3	ω	1	1	1	1

	χ_1	(χ_2, χ_3)	(χ_4, χ_5)	(χ_6, χ_7, χ_8)	$(\chi_9, \chi_{10}, \chi_{11})$	$(\chi_{12}, \chi_{13}, \chi_{14})$
SU(5)	1	1	1	1	1	1
S_4	1_1	2	2	3_1	3_1	3_1
Z_4	ω^2	ω^2	1	ω^3	1	ω

Table 1: Assignments of SU(5), S_4 , and Z_4 representations, where the phase factor ω is i .

Let us present the model of the quark and lepton flavor with the S_4 group in SU(5) GUT. In SU(5), matter fields are unified into a $10(q_1, u^c, e^c)_L$ and a $\bar{5}(d^c, l_e)_L$ dimensional representations. Three generations of $\bar{5}$, which are denoted by F_i , are assigned by 3_1 of S_4 . On the other hand, the third generation of the 10-dimensional representation is assigned by 1_1 of S_4 , so that the top quark Yukawa coupling is allowed in tree level. While, the first and the second generations are assigned 2 of S_4 . These 10-dimensional representations are denoted by T_3 and (T_1, T_2) , respectively. Right-handed neutrinos, which are SU(5) gauge singlets, are also assigned 1_1 and 2 for N_τ^c and (N_e^c, N_μ^c) , respectively ¹.

We introduce new scalars χ_i in addition to the 5-dimensional and $\bar{5}$ -dimensional Higgs of the SU(5), H_5 and $H_{\bar{5}}$ which are assigned 1_1 of S_4 . These new scalars are supposed to be SU(5) gauge singlets. The χ_1 scalar is assigned 1_1 , (χ_2, χ_3) , and (χ_4, χ_5) are 2, (χ_6, χ_7, χ_8) , $(\chi_9, \chi_{10}, \chi_{11})$, and $(\chi_{12}, \chi_{13}, \chi_{14})$ are 3_1 of the S_4 representations, respectively. The χ_1 and (χ_2, χ_3) scalars are coupled with the up type quark sector, (χ_4, χ_5) are coupled with the right-handed Majorana neutrino sector, (χ_6, χ_7, χ_8) are coupled with the Dirac neutrino sector, $(\chi_9, \chi_{10}, \chi_{11})$ and $(\chi_{12}, \chi_{13}, \chi_{14})$ are coupled with the charged lepton and down type quark sector, respectively. We also add Z_4 symmetry in order to obtain relevant couplings. The particle assignments of SU(5), S_4 , and Z_4 are summarized Table 1.

We can now write down the superpotential at the leading order in terms of the cut off scale Λ , which is taken to be the Planck scale. The SU(5) invariant superpotential of the

¹The similar assignments of right-handed neutrinos were presented in the first version of Ref.[49].

Yukawa sector respecting S_4 and Z_4 symmetries is given as

$$\begin{aligned}
w_{\text{SU}(5)}^{(0)} = & y_1^u(T_1, T_2) \otimes (T_1, T_2) \otimes \chi_1 \otimes H_5/\Lambda + y_2^u(T_1, T_2) \otimes (T_1, T_2) \otimes (\chi_2, \chi_3) \otimes H_5/\Lambda \\
& + y_3^u T_3 \otimes T_3 \otimes H_5 + M_1(N_e^c, N_\mu^c) \otimes (N_e^c, N_\mu^c) + M_2 N_\tau^c \otimes N_\tau^c \\
& + y^N(N_e^c, N_\mu^c) \otimes (N_e^c, N_\mu^c) \otimes (\chi_4, \chi_5) \\
& + y_1^D(N_e^c, N_\mu^c) \otimes (F_1, F_2, F_3) \otimes (\chi_6, \chi_7, \chi_8) \otimes H_5/\Lambda \\
& + y_2^D N_\tau^c \otimes (F_1, F_2, F_3) \otimes (\chi_6, \chi_7, \chi_8) \otimes H_5/\Lambda \\
& + y_1(F_1, F_2, F_3) \otimes (T_1, T_2) \otimes (\chi_9, \chi_{10}, \chi_{11}) \otimes H_5/\Lambda \\
& + y_2(F_1, F_2, F_3) \otimes T_3 \otimes (\chi_{12}, \chi_{13}, \chi_{14}) \otimes H_5/\Lambda,
\end{aligned} \tag{1}$$

where M_1 and M_2 are mass parameters for right-handed Majorana neutrinos, and Yukawa coupling constants y_i^a and y_i are complex in general. By decomposing this superpotential into the quark sector and the lepton sector, we can discuss mass matrices of quarks and leptons in following sections.

3 Lepton sector

At first, we begin to discuss the lepton sector of the superpotential $w_{\text{SU}(5)}^{(0)}$. Denoting Higgs doublets as h_u and h_d , the superpotential of the Yukawa sector respecting the $S_4 \times Z_4$ symmetry is given for charged leptons as

$$\begin{aligned}
w_l = & y_1 \left[\frac{e^c}{\sqrt{2}}(l_\mu \chi_{10} - l_\tau \chi_{11}) + \frac{\mu^c}{\sqrt{6}}(-2l_e \chi_9 + l_\mu \chi_{10} + l_\tau \chi_{11}) \right] h_d/\Lambda \\
& + y_2 \tau^c(l_e \chi_{12} + l_\mu \chi_{13} + l_\tau \chi_{14}) h_d/\Lambda.
\end{aligned} \tag{2}$$

For right-handed Majorana neutrinos, the superpotential is given as

$$\begin{aligned}
w_N = & M_1(N_e^c N_e^c + N_\mu^c N_\mu^c) + M_2 N_\tau^c N_\tau^c \\
& + y^N [(N_e^c N_\mu^c + N_\mu^c N_e^c) \chi_4 + (N_e^c N_e^c - N_\mu^c N_\mu^c) \chi_5],
\end{aligned} \tag{3}$$

and for neutrino Yukawa couplings, the superpotential is

$$\begin{aligned}
w_D = & y_1^D \left[\frac{N_e^c}{\sqrt{2}}(l_\mu \chi_7 - l_\tau \chi_8) + \frac{N_\mu^c}{\sqrt{6}}(-2l_e \chi_6 + l_\mu \chi_7 + l_\tau \chi_8) \right] h_u/\Lambda \\
& + y_2^D N_\tau^c(l_e \chi_6 + l_\mu \chi_7 + l_\tau \chi_8) h_u/\Lambda.
\end{aligned} \tag{4}$$

Higgs doublets h_u, h_d and gauge singlet scalars χ_i , are assumed to develop their vacuum expectation values (VEVs) as follows:

$$\begin{aligned}
\langle h_u \rangle = v_u, \quad \langle h_d \rangle = v_d, \quad \langle (\chi_4, \chi_5) \rangle = (u_4, u_5), \quad \langle (\chi_6, \chi_7, \chi_8) \rangle = (u_6, u_7, u_8), \\
\langle (\chi_9, \chi_{10}, \chi_{11}) \rangle = (u_9, u_{10}, u_{11}), \quad \langle (\chi_{12}, \chi_{13}, \chi_{14}) \rangle = (u_{12}, u_{13}, u_{14}),
\end{aligned} \tag{5}$$

which are supposed to be real. Then, we obtain the mass matrix for charged leptons as

$$M_l = y_1 v_d \begin{pmatrix} 0 & \alpha_{10}/\sqrt{2} & -\alpha_{11}/\sqrt{2} \\ -2\alpha_9/\sqrt{6} & \alpha_{10}/\sqrt{6} & \alpha_{11}/\sqrt{6} \\ 0 & 0 & 0 \end{pmatrix} + y_2 v_d \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \alpha_{12} & \alpha_{13} & \alpha_{14} \end{pmatrix}, \quad (6)$$

while the right-handed Majorana neutrino mass matrix is given as

$$M_N = \begin{pmatrix} M_1 + y^N \alpha_5 \Lambda & y^N \alpha_4 \Lambda & 0 \\ y^N \alpha_4 \Lambda & M_1 - y^N \alpha_5 \Lambda & 0 \\ 0 & 0 & M_2 \end{pmatrix}, \quad (7)$$

and the Dirac mass matrix of neutrinos is

$$M_D = y_1^D v_u \begin{pmatrix} 0 & \alpha_7/\sqrt{2} & -\alpha_8/\sqrt{2} \\ -2\alpha_6/\sqrt{6} & \alpha_7/\sqrt{6} & \alpha_8/\sqrt{6} \\ 0 & 0 & 0 \end{pmatrix} + y_2^D v_u \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \alpha_6 & \alpha_7 & \alpha_8 \end{pmatrix}, \quad (8)$$

where we denote $\alpha_i \equiv u_i/\Lambda$.

Let us discuss lepton masses and mixing angles by considering mass matrices in Eqs.(6), (7) and (8). In order to get the left-handed mixing of charged leptons, we investigate $M_l^\dagger M_l$:

$$M_l^\dagger M_l = v_d^2 \times \begin{pmatrix} \frac{2}{3}|y_1|^2 \alpha_9^2 + |y_2|^2 \alpha_{12}^2 & -\frac{1}{3}|y_1|^2 \alpha_9 \alpha_{10} + |y_2|^2 \alpha_{12} \alpha_{13} & -\frac{1}{3}|y_1|^2 \alpha_9 \alpha_{11} + |y_2|^2 \alpha_{12} \alpha_{14} \\ -\frac{1}{3}|y_1|^2 \alpha_9 \alpha_{10} + |y_2|^2 \alpha_{12} \alpha_{13} & \frac{2}{3}|y_1|^2 \alpha_{10}^2 + |y_2|^2 \alpha_{13}^2 & -\frac{1}{3}|y_1|^2 \alpha_{10} \alpha_{11} + |y_2|^2 \alpha_{13} \alpha_{14} \\ -\frac{1}{3}|y_1|^2 \alpha_9 \alpha_{11} + |y_2|^2 \alpha_{12} \alpha_{14} & -\frac{1}{3}|y_1|^2 \alpha_{10} \alpha_{11} + |y_2|^2 \alpha_{13} \alpha_{14} & \frac{2}{3}|y_1|^2 \alpha_{11}^2 + |y_2|^2 \alpha_{14}^2 \end{pmatrix}. \quad (9)$$

If we can take vacuum alignments $(u_9, u_{10}, u_{11}) = (u_9, u_{10}, 0)$ and $(u_{12}, u_{13}, u_{14}) = (0, 0, u_{14})$, that is $\alpha_{11} = \alpha_{12} = \alpha_{13} = 0$, we obtain

$$M_l^\dagger M_l = v_d^2 \begin{pmatrix} \frac{2}{3}|y_1|^2 \alpha_9^2 & -\frac{1}{3}|y_1|^2 \alpha_9 \alpha_{10} & 0 \\ -\frac{1}{3}|y_1|^2 \alpha_9 \alpha_{10} & \frac{2}{3}|y_1|^2 \alpha_{10}^2 & 0 \\ 0 & 0 & |y_2|^2 \alpha_{14}^2 \end{pmatrix}, \quad (10)$$

which gives $\theta_{13}^l = \theta_{23}^l = 0$, where θ_{ij}^l denote left-handed mixing angles to diagonalize the charged lepton mass matrix. Since the electron mass is tiny compared with the muon mass, we expect $\alpha_9 \ll \alpha_{10}$ and then we get the mixing angle θ_{12}^l as,

$$\tan \theta_{12}^l \approx -\frac{\alpha_9}{2\alpha_{10}}, \quad (11)$$

and charged lepton masses,

$$m_e^2 \approx \frac{1}{2}|y_1|^2 \alpha_9^2 v_d^2, \quad m_\mu^2 \approx \frac{2}{3}|y_1|^2 \alpha_{10}^2 v_d^2 + \frac{1}{6}|y_1|^2 \alpha_9^2 v_d^2 \approx \frac{2}{3}|y_1|^2 \alpha_{10}^2 v_d^2, \quad m_\tau^2 = |y_2|^2 \alpha_{14}^2 v_d^2. \quad (12)$$

Therefore, the mixing of θ_{12}^l is estimated as

$$|\tan \theta_{12}^l| \approx \frac{1}{\sqrt{3}} \frac{m_e}{m_\mu} \approx 2.8 \times 10^{-3}, \quad (13)$$

which is negligibly small. The mixing angle θ_{13}^l is at most $\mathcal{O}(m_e/m_\tau)$ even if we take into account of non-zero α_{12} . These tiny θ_{12}^l and θ_{13}^l hardly affect the magnitude of the lepton mixing matrix element U_{e3} , which will be discussed later.

It is noticed that one can take at the leading order the vacuum alignment $(u_9, u_{10}, u_{11}) = (0, u_{10}, 0)$ in order to guarantee $\alpha_9 \ll \alpha_{10}$, in which the electron mass vanishes. In conclusion, we find the charged lepton mass matrix to be almost diagonal one.

Taking vacuum alignments $(u_4, u_5) = (0, u_5)$ and $(u_6, u_7, u_8) = (u_6, u_6, u_6)$ in Eq.(7), the Majorana mass matrix of neutrinos turns to

$$M_N = \begin{pmatrix} M_1 + y^N \alpha_5 \Lambda & 0 & 0 \\ 0 & M_1 - y^N \alpha_5 \Lambda & 0 \\ 0 & 0 & M_2 \end{pmatrix}, \quad (14)$$

and the Dirac mass matrix of neutrinos turns to

$$M_D = y_1^D v_u \begin{pmatrix} 0 & \alpha_6/\sqrt{2} & -\alpha_6/\sqrt{2} \\ -2\alpha_6/\sqrt{6} & \alpha_6/\sqrt{6} & \alpha_6/\sqrt{6} \\ 0 & 0 & 0 \end{pmatrix} + y_2^D v_u \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \alpha_6 & \alpha_6 & \alpha_6 \end{pmatrix}. \quad (15)$$

By using the seesaw mechanism $M_\nu = M_D^T M_N^{-1} M_D$, the left-handed Majorana neutrino mass matrix is written as

$$M_\nu = \begin{pmatrix} a + \frac{2}{3}b & a - \frac{1}{3}b & a - \frac{1}{3}b \\ a - \frac{1}{3}b & a + \frac{1}{6}b + \frac{1}{2}c & a + \frac{1}{6}b - \frac{1}{2}c \\ a - \frac{1}{3}b & a + \frac{1}{6}b - \frac{1}{2}c & a + \frac{1}{6}b + \frac{1}{2}c \end{pmatrix}, \quad (16)$$

where

$$a = \frac{(y_2^D \alpha_6 v_u)^2}{M_2}, \quad b = \frac{(y_1^D \alpha_6 v_u)^2}{M_1 - y^N \alpha_5 \Lambda}, \quad c = \frac{(y_1^D \alpha_6 v_u)^2}{M_1 + y^N \alpha_5 \Lambda}. \quad (17)$$

The neutrino mass matrix is decomposed as

$$M_\nu = \frac{b+c}{2} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} + \frac{3a-b}{3} \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} + \frac{b-c}{2} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}. \quad (18)$$

As well known, the neutrino mass matrix with the tri-bimaximal mixing is expressed in terms of neutrino mass eigenvalues m_1 , m_2 and m_3 as

$$M_\nu = \frac{m_1 + m_3}{2} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} + \frac{m_2 - m_1}{3} \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} + \frac{m_1 - m_3}{2} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}. \quad (19)$$

Therefore, our neutrino mass matrix M_ν gives the tri-bimaximal mixing matrix $U_{\text{tri-bi}}$ and mass eigenvalues as follows:

$$U_{\text{tri-bi}} = \begin{pmatrix} \frac{2}{\sqrt{6}} & \frac{1}{\sqrt{3}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \end{pmatrix}, \quad m_1 = b, \quad m_2 = 3a, \quad m_3 = c. \quad (20)$$

We remind ourselves that the flavor mixing from the charged lepton sector is negligibly small.

Defining parameters $\mu_0 = v_u/\Lambda$, $\lambda_1 = M_1/\Lambda$ and $\lambda_2 = M_2/\Lambda$, and taking $y_1^D = y_2^D$, the observed values Δm_{atm}^2 and Δm_{sol}^2 are expressed as

$$\Delta m_{\text{atm}}^2 = -\frac{4y^N\alpha_5\lambda_1}{(\lambda_1^2 - y^{N^2}\alpha_5^2)^2} (y_1^D\alpha_6)^4\mu_0^2v_u^2, \quad \Delta m_{\text{sol}}^2 = \frac{9(\lambda_1^2 - y^N\alpha_5)^2 - \lambda_2^2}{\lambda_2^2(\lambda_1 - y^N\alpha_5)^2} (y_1^D\alpha_6)^4\mu_0^2v_u^2. \quad (21)$$

Putting $\Lambda = 2.43 \times 10^{18} \text{ GeV}$ and experimental values of $\Delta m_{\text{atm}}^2 = (2.1 - 2.8) \times 10^{-3} \text{ eV}^2$ and $\Delta m_{\text{sol}}^2 = (7.1 - 8.3) \times 10^{-5} \text{ eV}^2$ [1], we can estimate magnitudes of α_5 and α_6 in the case of the normal neutrino mass hierarchy. We show the numerical result in Figure 1, where we fix $y^N = -1$ and $y_1^D = 1$. We find $\alpha_6 \geq 0.1$, which is much larger than $\alpha_5 \approx 10^{-4} - 10^{-2}$.

Let us discuss a possible case of the non-vanishing U_{e3} , which is the deviation from the tri-bimaximal mixing. If $\alpha_4 \neq 0$, which corresponds to the non-vanishing u_4 , the left-handed Majorana neutrino mass matrix deviates from Eq.(16). After rotating by the tri-bimaximal mixing matrix ($\hat{M}_\nu = U_{\text{tri-bi}}^T M_\nu U_{\text{tri-bi}}$), we obtain off-diagonal elements in the neutrino mass matrix due to the non-zero α_4 as follows:

$$\hat{M}_\nu = \alpha_6^2 v_u^2 \begin{pmatrix} \frac{y_1^{D^2}(M_1 + y^N\alpha_5\Lambda)}{M_1^2 - y^{N^2}(\alpha_4^2 + \alpha_5^2)\Lambda^2} & 0 & -\frac{y_1^{D^2}y^N\alpha_4\Lambda}{M_1^2 - y^{N^2}(\alpha_4^2 + \alpha_5^2)\Lambda^2} \\ 0 & \frac{3y_2^{D^2}}{M_2} & 0 \\ -\frac{y_1^{D^2}y^N\alpha_4\Lambda}{M_1^2 - y^{N^2}(\alpha_4^2 + \alpha_5^2)\Lambda^2} & 0 & \frac{y_1^{D^2}(M_1 - y^N\alpha_5\Lambda)}{M_1^2 - y^{N^2}(\alpha_4^2 + \alpha_5^2)\Lambda^2} \end{pmatrix}. \quad (22)$$

Then the mixing angle $\delta\theta_{13}^\nu$, which diagonalizes this mass matrix, is given as

$$\tan 2\delta\theta_{13}^\nu = \frac{\alpha_4}{\alpha_5}, \quad (23)$$

which leads to

$$|U_{e3}| = \frac{2}{\sqrt{6}}|\delta\theta_{13}^\nu| \approx \frac{1}{\sqrt{6}}\left|\frac{\alpha_4}{\alpha_5}\right|, \quad |U_{e2}| = \frac{1}{\sqrt{3}}, \quad |U_{\mu 3}| \approx \left|\frac{1}{\sqrt{2}} + \frac{1}{2\sqrt{6}}\frac{\alpha_4}{\alpha_5}\right|. \quad (24)$$

Thus, the magnitude of U_{e3} is determined by the non-vanishing ratio α_4/α_5 .

4 Quark sector

In this section, we discuss quark sector of the superpotential $w_{\text{SU}(5)}^{(0)}$. For up type quarks, the superpotential of the Yukawa sector with $S_4 \times Z_4$ is given as

$$w_u = y_1^u(u^c q_1 + c^c q_2)\chi_1 h_u/\Lambda + y_2^u[(u^c q_2 + c^c q_1)\chi_2 + (u^c q_1 - c^c q_2)\chi_3] h_u/\Lambda + y_3^u t^c q_3 h_u. \quad (25)$$




Figure 1: The allowed region on the $\alpha_5 - \alpha_6$ plane.

For down type quarks, we can write the superpotential as follows:

$$w_d = y_1 \left[\frac{1}{\sqrt{2}}(s^c \chi_{10} - b^c \chi_{11})q_1 + \frac{1}{\sqrt{6}}(-2d^c \chi_9 + s^c \chi_{10} + b^c \chi_{11})q_2 \right] h_d/\Lambda \\ + y_2(d^c \chi_{12} + s^c \chi_{13} + b^c \chi_{14})q_3 h_d/\Lambda. \quad (26)$$

We assume that scalar fields, χ_i , develop their VEVs as follows:

$$\langle \chi_1 \rangle = u_1, \quad \langle (\chi_2, \chi_3) \rangle = (u_2, u_3), \\ \langle (\chi_9, \chi_{10}, \chi_{11}) \rangle = (u_9, u_{10}, u_{11}), \quad \langle (\chi_{12}, \chi_{13}, \chi_{14}) \rangle = (u_{12}, u_{13}, u_{14}). \quad (27)$$

Then, the mass matrix for up type quarks is given as

$$M_u = v_u \begin{pmatrix} y_1^u \alpha_1 + y_2^u \alpha_3 & y_2^u \alpha_2 & 0 \\ y_2^u \alpha_2 & y_1^u \alpha_1 - y_2^u \alpha_3 & 0 \\ 0 & 0 & y_3^u \end{pmatrix}, \quad (28)$$

and the down type quark mass matrix is given as

$$M_d = y_1 v_d \begin{pmatrix} 0 & -2\alpha_9/\sqrt{6} & 0 \\ \alpha_{10}/\sqrt{2} & \alpha_{10}/\sqrt{6} & 0 \\ -\alpha_{11}/\sqrt{2} & \alpha_{11}/\sqrt{6} & 0 \end{pmatrix} + y_2 v_d \begin{pmatrix} 0 & 0 & \alpha_{12} \\ 0 & 0 & \alpha_{13} \\ 0 & 0 & \alpha_{14} \end{pmatrix}. \quad (29)$$

Let us discuss masses and mixing of the quark sector. For up type quarks, if we take

$$\alpha_3 = 0, \quad y_1^u \alpha_1 = y_2^u \alpha_2, \quad (30)$$

which will be reexamined to get observed CKM mixing angles in section 5.2, then, we have

$$M_u = v_u \begin{pmatrix} y_1^u \alpha_1 & y_1^u \alpha_1 & 0 \\ y_1^u \alpha_1 & y_1^u \alpha_1 & 0 \\ 0 & 0 & y_3^u \end{pmatrix}, \quad (31)$$

which is diagonalized by the orthogonal matrix U_u

$$U_u = \begin{pmatrix} \cos 45^\circ & \sin 45^\circ & 0 \\ -\sin 45^\circ & \cos 45^\circ & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (32)$$

The up type quark masses are given as

$$m_u = 0, \quad m_c = 2y_1^u v_u \alpha_1, \quad m_t = y_3^u v_u. \quad (33)$$

For down type quarks, putting $\alpha_{11} = \alpha_{12} = \alpha_{13} = 0$, which is the condition in the charged lepton sector, we have

$$M_d^\dagger M_d = v_d^2 \begin{pmatrix} \frac{1}{2}|y_1|^2 \alpha_{10}^2 & \frac{1}{2\sqrt{3}}|y_1|^2 \alpha_{10}^2 & 0 \\ \frac{1}{2\sqrt{3}}|y_1|^2 \alpha_{10}^2 & \frac{1}{6}|y_1|^2 (4\alpha_9^2 + \alpha_{10}^2) & 0 \\ 0 & 0 & |y_2|^2 \alpha_{14}^2 \end{pmatrix}. \quad (34)$$

Then, the mass matrix is diagonalized by the orthogonal matrix U_d as

$$U_d = \begin{pmatrix} \cos 60^\circ & \sin 60^\circ & 0 \\ -\sin 60^\circ & \cos 60^\circ & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (35)$$

where the small α_9 is neglected. The down type quark masses are given as

$$m_d^2 \approx \frac{1}{2}|y_1|^2 \alpha_9^2 v_d^2, \quad m_s^2 \approx \frac{2}{3}|y_1|^2 \alpha_{10}^2 v_d^2, \quad m_b^2 \approx |y_2|^2 \alpha_{14}^2 v_d^2, \quad (36)$$

which are the same ones of charged lepton masses in Eq.(12).

Now, we get the CKM matrix as follows:

$$V^{CKM} = U_u^\dagger U_d = \begin{pmatrix} \cos 15^\circ & \sin 15^\circ & 0 \\ -\sin 15^\circ & \cos 15^\circ & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (37)$$

Therefore, in our prototype model of SU(5) GUT with the S_4 flavor symmetry, the quark sector has a non-vanishing mixing angle 15° only between the first and second generations while the lepton flavor mixing is tri-bimaximal. In order to get the non-vanishing but small mixing angles V_{cb}^{CKM} and V_{ub}^{CKM} , we improve the prototype model in the next section.

5 Improved S_4 flavor model in SU(5) GUT

We improve the prototype model to get the observed quark and lepton mass spectra and the CKM mixing matrix. We introduce the SU(5) 45-dimensional Higgs h_{45} , which is required to get the difference between the charged lepton mass spectrum and the down type quark mass spectrum. Moreover, we add a S_4 doublet (χ'_2, χ'_3) and a S_4 triplet $(\chi'_9, \chi'_{10}, \chi'_{11})$, which are SU(5) gauge singlet scalars. These assignments of SU(5), S_4 , and Z_4 are summarized Table 2. Since the additional scalars do not contribute to the neutrino sector, the result of the neutrino sector in the prototype model is not changed. Therefore, we discuss only the charged lepton sector and the quark sector in this section.

	h_{45}	(χ'_2, χ'_3)	$(\chi'_9, \chi'_{10}, \chi'_{11})$
SU(5)	45	1	1
S_4	1_1	2	3_1
Z_4	ω^2	ω^3	ω^2

Table 2: Assignments of additional scalars in SU(5), S_4 , and Z_4 representations.

The superpotential of the Yukawa sector respecting the SU(5), S_4 and Z_4 symmetries is given as

$$w_{\text{SU}(5)} = w_{\text{SU}(5)}^{(0)} + w_{\text{SU}(5)}^{(1)}, \quad (38)$$

where we denote

$$\begin{aligned} w_{\text{SU}(5)}^{(1)} &= y_4^u(T_1, T_2) \otimes T_3 \otimes (\chi'_2, \chi'_3) \otimes H_5 \\ &+ y'_1(F_1, F_2, F_3) \otimes (T_1, T_2) \otimes (\chi'_9, \chi'_{10}, \chi'_{11}) \otimes h_{45}. \end{aligned} \quad (39)$$

5.1 Improved lepton sector

Let us discuss the improved lepton sector of the superpotential $w_{\text{SU}(5)}$. The superpotential of the charged lepton sector with $S_4 \times Z_4$ is given as

$$\begin{aligned} w_l = & y_1 \left[\frac{e^c}{\sqrt{2}}(l_\mu \chi_{10} - l_\tau \chi_{11}) + \frac{\mu^c}{\sqrt{6}}(-2l_e \chi_9 + l_\mu \chi_{10} + l_\tau \chi_{11}) \right] h_d/\Lambda \\ & - 3y'_1 \left[\frac{e^c}{\sqrt{2}}(l_\mu \chi'_{10} - l_\tau \chi'_{11}) + \frac{\mu^c}{\sqrt{6}}(-2l_e \chi'_9 + l_\mu \chi'_{10} + l_\tau \chi'_{11}) \right] h_{45}/\Lambda \\ & + y_2 \tau^c (l_e \chi_{12} + l_\mu \chi_{13} + l_\tau \chi_{14}) h_d/\Lambda. \end{aligned} \quad (40)$$

We denote their VEVs as follows:

$$\begin{aligned} \langle h_{45} \rangle &= v_{45}, \quad \langle (\chi_9, \chi_{10}, \chi_{11}) \rangle = (u_9, u_{10}, u_{11}), \\ \langle (\chi'_9, \chi'_{10}, \chi'_{11}) \rangle &= (u'_9, u'_{10}, u'_{11}), \quad \langle (\chi_{12}, \chi_{13}, \chi_{14}) \rangle = (u_{12}, u_{13}, u_{14}). \end{aligned} \quad (41)$$

Then, we obtain the mass matrix for charged leptons:

$$\begin{aligned} M_l = & y_1 v_d \begin{pmatrix} 0 & \alpha_{10}/\sqrt{2} & -\alpha_{11}/\sqrt{2} \\ -2\alpha_9/\sqrt{6} & \alpha_{10}/\sqrt{6} & \alpha_{11}/\sqrt{6} \\ 0 & 0 & 0 \end{pmatrix} + y_2 v_d \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \alpha_{12} & \alpha_{13} & \alpha_{14} \end{pmatrix} \\ & - 3y'_1 v_{45} \begin{pmatrix} 0 & \alpha'_{10}/\sqrt{2} & -\alpha'_{11}/\sqrt{2} \\ -2\alpha'_9/\sqrt{6} & \alpha'_{10}/\sqrt{6} & \alpha'_{11}/\sqrt{6} \\ 0 & 0 & 0 \end{pmatrix}, \end{aligned} \quad (42)$$

where we denote $\alpha_i = u_i/\Lambda$ and $\alpha'_j = u'_j/\Lambda$. It is noticed that the third matrix in the right hand side is the additional one compared with the mass matrix of the prototype model in Eq.(6).

Masses and mixing angles of the charged lepton sector are similar to those of the prototype model in Eqs.(12) and (13). If we can take the vacuum alignments $(u_9, u_{10}, u_{11}) = (u_9, u_{10}, 0)$, $(u'_9, u'_{10}, u'_{11}) = (u'_9, u'_{10}, 0)$ and $(u_{12}, u_{13}, u_{14}) = (0, 0, u_{14})$, that is $\alpha_{11} = \alpha'_{11} = \alpha_{12} = \alpha_{13} = 0$, we obtain charged lepton mass matrix as follow:

$$M_l = v_d \begin{pmatrix} 0 & (y_1 \alpha_{10} - 3\bar{y}_1 \alpha'_{10})/\sqrt{2} & 0 \\ -2(y_1 \alpha_9 - 3\bar{y}_1 \alpha'_9)/\sqrt{6} & (y_1 \alpha_{10} - 3\bar{y}_1 \alpha'_{10})/\sqrt{6} & 0 \\ 0 & 0 & y_2 \alpha_{14} \end{pmatrix}, \quad (43)$$

where we replace $y'_1 v_{45}$ with $\bar{y}_1 v_d$. Since we have

$$\begin{aligned} M_l^\dagger M_l = & v_d^2 \times \\ & \begin{pmatrix} \frac{2}{3}|y_1 \alpha_9 - 3\bar{y}_1 \alpha'_9|^2 & -\frac{1}{3}(y_1^* \alpha_9 - 3\bar{y}_1^* \alpha'_9)(y_1 \alpha_{10} - 3\bar{y}_1 \alpha'_{10}) & 0 \\ -\frac{1}{3}(y_1 \alpha_9 - 3\bar{y}_1 \alpha'_9)(y_1^* \alpha_{10} - 3\bar{y}_1^* \alpha'_{10}) & \frac{2}{3}|y_1 \alpha_{10} - 3\bar{y}_1 \alpha'_{10}|^2 & 0 \\ 0 & 0 & |y_2|^2 \alpha_{14}^2 \end{pmatrix}, \end{aligned} \quad (44)$$

masses and mixing angles of the charged leptons as follows:

$$m_e^2 \approx \frac{1}{2}|y_1\alpha_9 - 3\bar{y}_1\alpha'_9|^2 v_d^2, \quad m_\mu^2 \approx \frac{2}{3}|y_1\alpha_{10} - 3\bar{y}_1\alpha'_{10}|^2 v_d^2, \quad m_\tau^2 \approx |y_2|^2 \alpha_{14}^2 v_d^2, \\ |\theta_{12}^l| = \left| -\frac{y_1\alpha_9 - 3\bar{y}_1\alpha'_9}{2(y_1\alpha_{10} - 3\bar{y}_1\alpha'_{10})} \right| \approx \frac{1}{\sqrt{3}} \frac{m_e}{m_\mu} \approx 2.8 \times 10^{-3}, \quad \theta_{23}^l = 0, \quad \theta_{13}^l = 0. \quad (45)$$

Thus, the charged lepton mass matrix is almost diagonal, and so the tri-bimaximal mixing of neutrino flavors is also realized in this improved model.

5.2 Improved quark sector

Let us discuss the quark sector of the superpotential $w_{\text{SU}(5)}$. For up type quarks, the superpotential respecting $S_4 \times Z_4$ is given as

$$w_u = y_1^u (u^c q_1 + c^c q_2) \chi_1 h_u / \Lambda \\ + y_2^u [(u^c q_2 + c^c q_1) \chi_2 + (u^c q_1 - c^c q_2) \chi_3] h_u / \Lambda \\ + y_3^u t^c q_3 h_u \\ + y_4^u [(u^c \chi'_2 + c^c \chi'_3) q_3 + t^c (q_1 \chi'_2 + q_2 \chi'_3)] h_u / \Lambda. \quad (46)$$

For down type quarks, we can write the superpotential as follows:

$$w_d = y_1 \left[\frac{1}{\sqrt{2}} (s^c \chi_{10} - b^c \chi_{11}) q_1 + \frac{1}{\sqrt{6}} (-2d^c \chi_9 + s^c \chi_{10} + b^c \chi_{11}) q_2 \right] h_d / \Lambda \\ + y'_1 \left[\frac{1}{\sqrt{2}} (s^c \chi'_{10} - b^c \chi'_{11}) q_1 + \frac{1}{\sqrt{6}} (-2d^c \chi'_9 + s^c \chi'_{10} + b^c \chi'_{11}) q_2 \right] v_{45} / \Lambda \\ + y_2 (d^c \chi_{12} + s^c \chi_{13} + b^c \chi_{14}) q_3 h_d / \Lambda. \quad (47)$$

We denote their VEVs as follows:

$$\langle \chi_1 \rangle = u_1, \quad \langle (\chi_2, \chi_3) \rangle = (u_2, u_3), \quad \langle (\chi'_2, \chi'_3) \rangle = (u'_2, u'_3), \\ \langle (\chi_9, \chi_{10}, \chi_{11}) \rangle = (u_9, u_{10}, u_{11}), \quad \langle (\chi'_9, \chi'_{10}, \chi'_{11}) \rangle = (u'_9, u'_{10}, u'_{11}), \\ \langle (\chi_{12}, \chi_{13}, \chi_{14}) \rangle = (u_{12}, u_{13}, u_{14}). \quad (48)$$

Then, we obtain the mass matrix for up type quarks is given as

$$M_u = v_u \begin{pmatrix} y_1^u \alpha_1 + y_2^u \alpha_3 & y_2^u \alpha_2 & y_4^u \alpha'_2 \\ y_2^u \alpha_2 & y_1^u \alpha_1 - y_2^u \alpha_3 & y_4^u \alpha'_3 \\ y_4^u \alpha'_2 & y_4^u \alpha'_3 & y_3^u \end{pmatrix}, \quad (49)$$

while the down type quark mass matrix is given as

$$M_d = y_1 v_d \begin{pmatrix} 0 & -2\alpha_9/\sqrt{6} & 0 \\ \alpha_{10}/\sqrt{2} & \alpha_{10}/\sqrt{6} & 0 \\ -\alpha_{11}/\sqrt{2} & \alpha_{11}/\sqrt{6} & 0 \end{pmatrix} + y_2 v_d \begin{pmatrix} 0 & 0 & \alpha_{12} \\ 0 & 0 & \alpha_{13} \\ 0 & 0 & \alpha_{14} \end{pmatrix} \\ + y'_1 v_{45} \begin{pmatrix} 0 & -2\alpha'_9/\sqrt{6} & 0 \\ \alpha'_{10}/\sqrt{2} & \alpha'_{10}/\sqrt{6} & 0 \\ -\alpha'_{11}/\sqrt{2} & \alpha'_{11}/\sqrt{6} & 0 \end{pmatrix}. \quad (50)$$

We consider the quark mixing. The up type quark mass matrix (49) turns to the following one after rotating by $\theta_{12}^u = 45^\circ$:

$$\hat{M}_u = v_u \begin{pmatrix} y_1^u \alpha_1 - y_2^u \alpha_2 & y_2^u \alpha_3 & \frac{y_4^u}{\sqrt{2}}(\alpha'_2 - \alpha'_3) \\ y_2^u \alpha_3 & y_1^u \alpha_1 + y_2^u \alpha_2 & \frac{y_4^u}{\sqrt{2}}(\alpha'_2 + \alpha'_3) \\ \frac{y_4^u}{\sqrt{2}}(\alpha'_2 - \alpha'_3) & \frac{y_4^u}{\sqrt{2}}(\alpha'_2 + \alpha'_3) & y_3^u \end{pmatrix}. \quad (51)$$

In order to obtain the non-vanishing quark mixing of V_{cb}^{CKM} and V_{ub}^{CKM} , we take

$$y_2^u \alpha_3 \gg y_1^u \alpha_1, y_2^u \alpha_2, \quad \alpha'_2 = \alpha'_3, \quad (52)$$

which are realized by vacuum alignments $u_1 = 0$ ², $(u_2, u_3) = (0, u_3)$ and $(u'_2, u'_3) = (u'_2, u'_2)$. This situation of VEVs is completely different from that of the prototype model as seen in Eq.(30), in which V_{cb}^{CKM} and V_{ub}^{CKM} vanish. Then, we obtain the so-called Fritzsch-type mass matrix [55]

$$\hat{M}_u \simeq v_u \begin{pmatrix} 0 & y_2^u \alpha_3 & 0 \\ y_2^u \alpha_3 & 0 & \sqrt{2} y_4^u \alpha'_2 \\ 0 & \sqrt{2} y_4^u \alpha'_2 & y_3^u \end{pmatrix}. \quad (53)$$

As well known, the complex phases in this 3×3 matrix can be removed by the phase matrix P as $P^\dagger \hat{M}_u P$;

$$P = \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{-i\rho} & 0 \\ 0 & 0 & e^{-i\sigma} \end{pmatrix}. \quad (54)$$

Therefore, up type quark masses are

$$m_u = \left| \frac{y_3^u y_2^{u2} \alpha_3^2}{2 y_4^{u2} \alpha_2'^2} \right| v_u, \quad m_c = \left| -\frac{2 y_4^{u2}}{y_3^u} \alpha_2'^2 \right| v_u, \quad m_t = |y_3^u| v_u, \quad (55)$$

and the mixing matrix to diagonalize $\hat{M}_u$ in Eq.(53), V_F ($M_u^{\text{diagonal}} = V_F^\dagger \hat{M}_u V_F$), is

$$V_F \approx \begin{pmatrix} 1 & \sqrt{\frac{m_u}{m_c}} & -\sqrt{\frac{m_u}{m_t}} \\ -\sqrt{\frac{m_u}{m_c}} & 1 & \sqrt{\frac{m_c}{m_t}} \\ \sqrt{\frac{m_u}{m_t}} & -\sqrt{\frac{m_c}{m_t}} & 1 \end{pmatrix}. \quad (56)$$

The conditions from the lepton sector $\alpha_{11} = \alpha'_{11} = \alpha_{12} = \alpha_{13} = 0$ give the down type quark mass matrix:

$$M_d = v_d \begin{pmatrix} 0 & -2(y_1 \alpha_9 + \bar{y}_1 \alpha'_9)/\sqrt{6} & 0 \\ (y_1 \alpha_{10} + \bar{y}_1 \alpha'_{10})/\sqrt{2} & (y_1 \alpha_{10} + \bar{y}_1 \alpha'_{10})/\sqrt{6} & 0 \\ 0 & 0 & y_2 \alpha_{14} \end{pmatrix}, \quad (57)$$

where we denote $\bar{y}_1 v_d = y'_1 v_{45}$. Then, we have

$$M_d^\dagger M_d = v_d^2 \begin{pmatrix} \frac{1}{2} |y_1 \alpha_{10} + \bar{y}_1 \alpha'_{10}|^2 & \frac{1}{2\sqrt{3}} |y_1 \alpha_{10} + \bar{y}_1 \alpha'_{10}|^2 & 0 \\ \frac{1}{2\sqrt{3}} |y_1 \alpha_{10} + \bar{y}_1 \alpha'_{10}|^2 & \frac{1}{6} (4|y_1 \alpha_9 + \bar{y}_1 \alpha'_9|^2 + |y_1 \alpha_{10} + \bar{y}_1 \alpha'_{10}|^2) & 0 \\ 0 & 0 & |y_2|^2 \alpha_{14}^2 \end{pmatrix}. \quad (58)$$

²One may consider to remove χ_1 , which is S_4 singlet, in our scheme.

After rotating by U_d in Eq.(35), $M_d^\dagger M_d$ turns to be

$$v_d^2 \begin{pmatrix} \frac{1}{2}|y_1\alpha_9 + \bar{y}_1\alpha'_9|^2 & -\frac{1}{2\sqrt{3}}|y_1\alpha_9 + \bar{y}_1\alpha'_9|^2 & 0 \\ -\frac{1}{2\sqrt{3}}|y_1\alpha_9 + \bar{y}_1\alpha'_9|^2 & \frac{1}{6}(|y_1\alpha_9 + \bar{y}_1\alpha'_9|^2 + 4|y_1\alpha_{10} + \bar{y}_1\alpha'_{10}|^2) & 0 \\ 0 & 0 & |y_2|^2\alpha_{14}^2 \end{pmatrix}. \quad (59)$$

Then, down type quark masses are given as

$$m_d^2 \approx \frac{1}{2}|y_1\alpha_9 + \bar{y}_1\alpha'_9|^2 v_d^2, \quad m_s^2 \approx \frac{2}{3}|y_1\alpha_{10} + \bar{y}_1\alpha'_{10}|^2 v_d^2, \quad m_b^2 \approx |y_2|^2 \alpha_{14}^2 v_d^2, \quad (60)$$

and the mixing angle θ_{12}^d is $60^\circ + \delta\theta_{12}^d$, where

$$\delta\theta_{12}^d = -\frac{\sqrt{3}|y_1\alpha_9 + \bar{y}_1\alpha'_9|^2}{4|y_1\alpha_{10} + \bar{y}_1\alpha'_{10}|^2} = -\frac{m_d^2}{\sqrt{3}m_s^2} \approx -1.5 \times 10^{-3}. \quad (61)$$

Therefore, θ_{12}^d is almost 60° . We add comments on θ_{23}^d and θ_{13}^d , which vanish in our scheme because of $\alpha_{11} = \alpha'_{11} = \alpha_{12} = \alpha_{13} = 0$. Non-vanishing α_{ij} lead to

$$\theta_{13}^d \approx -\frac{y_1\alpha_{11}}{\sqrt{2}y_2\alpha_{14}}, \quad \theta_{23}^d \approx \frac{2(y_1\alpha_{10} + \bar{y}_1\alpha'_{10})\alpha_{13} - y_1\alpha_{11}\alpha_{14}}{\sqrt{6}y_2\alpha_{14}^2}, \quad (62)$$

where the complex phases of Yukawa couplings are neglected. These mixing angles are expected to be tiny as far as $\alpha_{14} \gg \alpha_{11}, \alpha_{13}$.

Let us discuss the CKM matrix. The unitary matrices diagonalizing the up type quark mass matrix and the down type quark one, U_u and U_d are given, respectively,

$$U_u = \begin{pmatrix} \cos 45^\circ & \sin 45^\circ & 0 \\ -\sin 45^\circ & \cos 45^\circ & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{-i\rho} & 0 \\ 0 & 0 & e^{-i\sigma} \end{pmatrix} \begin{pmatrix} 1 & \sqrt{\frac{m_u}{m_c}} & -\sqrt{\frac{m_u}{m_t}} \\ -\sqrt{\frac{m_u}{m_c}} & 1 & \sqrt{\frac{m_c}{m_t}} \\ \sqrt{\frac{m_u}{m_t}} & -\sqrt{\frac{m_c}{m_t}} & 1 \end{pmatrix},$$

$$U_d = \begin{pmatrix} \cos 60^\circ & \sin 60^\circ & 0 \\ -\sin 60^\circ & \cos 60^\circ & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (63)$$

Therefore, the CKM matrix is written as

$$V^{CKM} = U_u^\dagger U_d = \begin{pmatrix} 1 & -\sqrt{\frac{m_u}{m_c}} & \sqrt{\frac{m_u}{m_t}} \\ \sqrt{\frac{m_u}{m_c}} & 1 & -\sqrt{\frac{m_c}{m_t}} \\ -\sqrt{\frac{m_u}{m_t}} & \sqrt{\frac{m_c}{m_t}} & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{i\rho} & 0 \\ 0 & 0 & e^{i\sigma} \end{pmatrix} \begin{pmatrix} \cos 15^\circ & \sin 15^\circ & 0 \\ -\sin 15^\circ & \cos 15^\circ & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (64)$$

The relevant CKM mixing elements are given as

$$|V_{us}^{CKM}| = \left| \sin 15^\circ - \cos 15^\circ \sqrt{\frac{m_u}{m_c}} e^{i\rho} \right|, \quad |V_{cb}^{CKM}| = \sqrt{\frac{m_c}{m_t}}, \quad |V_{ub}^{CKM}| = \sqrt{\frac{m_u}{m_t}}. \quad (65)$$

In the limit of neglecting the CP violating phase, $\rho = 0$, putting typical values at the GUT scale $m_u = 1.04 \times 10^{-3} \text{ GeV}$, $m_c = 302 \times 10^{-3} \text{ GeV}$, $m_t = 129 \text{ GeV}$, which are derived in Ref. [56], we predict

$$|V_{us}^{CKM}| = 0.202, \quad |V_{cb}^{CKM}| = 0.048, \quad |V_{ub}^{CKM}| = 0.003. \quad (66)$$

By adjusting the non-zero phase $\rho = 50^\circ$, we can get the central value of the observed Cabbibo angle 0.226. Another phase σ is still a free parameter. Our predicted V_{cb}^{CKM} and V_{ub}^{CKM} are somewhat different from the central values of observed mixing angles. If we take into account the down type quark mixing angles in Eq.(62), which are neglected in this stage, we expect to improve the situation including the phase σ . Then, we can discuss the CP violating phenomena in the CKM scheme by using two phases ρ and σ .

The difference between the charged lepton mass spectrum and the down type quark one is occurred by the 45-dimensional Higgs. The masses of charged leptons and down type quarks are given as:

$$\begin{aligned} m_e^2 &= \frac{1}{2}|y_1\alpha_9 - 3\bar{y}_1\alpha'_9|^2 v_d^2, & m_\mu^2 &= \frac{2}{3}|y_1\alpha_{10} - 3\bar{y}_1\alpha'_{10}|^2 v_d^2, & m_\tau^2 &= |y_2|^2 \alpha_{14}^2 v_d^2, \\ m_d^2 &= \frac{1}{2}|y_1\alpha_9 + \bar{y}_1\alpha'_9|^2 v_d^2, & m_s^2 &= \frac{2}{3}|y_1\alpha_{10} + \bar{y}_1\alpha'_{10}|^2 v_d^2, & m_b^2 &= |y_2|^2 \alpha_{14}^2 v_d^2. \end{aligned} \quad (67)$$

In order to get the power ratios of masses $m_e^2 : m_d^2 = 1 : 9$, $m_\mu^2 : m_s^2 = 9 : 1$, which is consistent with the observed mass spectra at the GUT scale, we require the following conditions:

$$\begin{aligned} y_1\alpha_9 &= 5\bar{y}_1\alpha'_9, & \text{or} & & y_1\alpha_9 &= 2\bar{y}_1\alpha'_9, \\ y_1\alpha_{10} &= -3\bar{y}_1\alpha'_{10}, & \text{or} & & y_1\alpha_{10} &= 0. \end{aligned} \quad (68)$$

6 Summary

We have presented a flavor model with the S_4 symmetry to unify quarks and leptons in the framework of the SU(5) GUT. Three generations of $\bar{5}$ -plets in SU(5) are assigned 3_1 of S_4 while the first and the second generations of 10-plets in SU(5) are assigned to be 2 of S_4 , and the third generation of 10-plet is to be 1_1 of S_4 . These assignments of S_4 for $\bar{5}$ and 10 lead to the completely different structure of quark and lepton mass matrices. Right-handed neutrinos, which are SU(5) gauge singlets, are also assigned 2 for the first and second generations and 1_1 for the third generation, respectively. These assignments are essential to realize the tri-bimaximal mixing of neutrino flavors. Vacuum alignments of scalars are also required to realize the tri-bimaximal mixing of neutrino flavors. Our model predicts the quark mixing as well as the tri-bimaximal mixing of leptons. Especially, the Cabbibo angle is predicted to be 15° in the limit of the vacuum alignment. We improve the model to predict observed CKM mixing angles as well as the non-vanishing U_{e3} of the neutrino flavor mixing. The deviation from 15° in $|V_{us}^{\text{CKM}}|$ is given by $\sqrt{m_u/m_c}$, while the non-vanishing $|V_{cb}^{\text{CKM}}|$ and $|V_{ub}^{\text{CKM}}|$ are given by $\sqrt{m_c/m_t}$ and $\sqrt{m_u/m_t}$, respectively. The non-vanishing U_{e3} of the neutrino flavor mixing is independent of these deviations. We will discuss the CP violating phenomena in the CKM scheme elsewhere by taking account of the corrections of the down type quark sector.

Acknowledgement

The work of M.T. has been supported by the Grant-in-Aid for Science Research of the Ministry of Education, Science, and Culture of Japan No. 17540243.

Appendix

A Multiplication rules of S_4

We present the relevant multiplication rules of S_4 . The two-dimensional one is given as

$$(a_1, a_2)_2 \times (b_1, b_2)_2 = (a_1b_1 + a_2b_2)_{1_1} + (-a_1b_2 + a_2b_1)_{1_2} + (a_1b_2 + a_2b_1, a_1b_1 - a_2b_2)_2. \quad (69)$$

For three-dimensional representation, the product is given as

$$\begin{aligned} (a_1, a_2, a_3)_{3_1} \times (b_1, b_2, b_3)_{3_1} = & (a_1b_1 + a_2b_2 + a_3b_3)_{1_1} \\ & + \left[\frac{1}{\sqrt{2}}(a_2b_2 - a_3b_3), \frac{1}{\sqrt{6}}(-2a_1b_1 + a_2b_2 + a_3b_3) \right]_{2_1} \\ & + (a_2b_3 + a_3b_2, a_1b_3 + a_3b_1, a_1b_2 + a_2b_1)_{3_1} \\ & + (a_3b_2 - a_2b_3, a_1b_3 - a_3b_1, a_2b_1 - a_1b_2)_{3_2}. \end{aligned} \quad (70)$$

The product of two- and three-dimensional representations is given as

$$\begin{aligned} (a_1, a_2)_2 \times (b_1, b_2, b_3)_{3_1} = & \left[a_2b_1, -\frac{1}{2}(\sqrt{3}a_1b_2 + a_2b_2), \frac{1}{2}(\sqrt{3}a_1b_3 - a_2b_3) \right]_{3_1} \\ & + \left[a_1b_1, \frac{1}{2}(\sqrt{3}a_2b_2 - a_1b_2), -\frac{1}{2}(\sqrt{3}a_2b_3 + a_1b_3) \right]_{3_2}. \end{aligned} \quad (71)$$

B Vacuum alignments and magnitudes of VEVs

In our model, we need vacuum alignments of scalar fields χ_i of S_4 doublets and triplets. Vacuum alignments are summarized at the leading order as follows:

$$\begin{aligned} \chi_1 = 0, \quad (\chi_2, \chi_3) = (0, 1), \quad (\chi'_2, \chi'_3) = (1, 1), \quad (\chi_4, \chi_5) = (0, 1), \quad (\chi_6, \chi_7, \chi_8) = (1, 1, 1) \\ (\chi_9, \chi_{10}, \chi_{11}) = (0, 1, 0), \quad (\chi'_9, \chi'_{10}, \chi'_{11}) = (0, 1, 0), \quad (\chi_{12}, \chi_{13}, \chi_{14}) = (0, 0, 1), \end{aligned} \quad (72)$$

where magnitudes are given in arbitrary units. Non-vanishing m_e and m_d require tiny deviations from zeros for χ_9 and χ'_9 , which could be realized in the next leading order.

In order to show magnitudes of VEVs, we estimate α_i and α'_j . These are given as follows:

$$\begin{aligned} \alpha_1 = \alpha_2 = \alpha_4 = \alpha_{11} = \alpha'_{11} = \alpha_{12} = \alpha_{13} = 0, \quad \alpha'_2 = \alpha'_3 = \sqrt{\left| \frac{y_3^u m_c}{2y_4^{u2} v_u} \right|}, \quad \alpha_3 = \sqrt{\frac{m_u m_c}{y_2^{u2} v_u^2}}, \\ \alpha_5 = \frac{(3y_2^{D2} m_1 - y_1^{D2} m_2) M_2}{3y_2^{D2} y^N m_1 \Lambda}, \quad \alpha_6 = \alpha_7 = \alpha_8 = \frac{\sqrt{m_2 M_2}}{\sqrt{3} y_2^D v_u}, \quad \alpha_9 = \frac{\sqrt{2}(3m_d + m_e)}{4y_1 v_d}, \\ \alpha'_9 = \frac{\sqrt{2}(m_d - m_e)}{4\bar{y}_1 v_d}, \quad \alpha_{10} = \frac{\sqrt{3}(3m_s + m_\mu)}{4\sqrt{2} y_1 v_d}, \quad \alpha'_{10} = \frac{\sqrt{3}(m_s - m_\mu)}{4\sqrt{2} \bar{y}_1 v_d}, \quad \alpha_{14} = \frac{m_b}{y_2 v_d}. \end{aligned} \quad (73)$$

Putting typical values of quark masses at the GUT scale [56], $M_2 = 10^{16} \text{GeV}$, and $\tan \beta = 3$ ($v_d \approx 55 \text{GeV}$, $v_u \approx 165 \text{GeV}$) with taking 1 for Yukawa couplings, we have

$$\begin{aligned} \alpha'_2 \sim 0.03, \quad \alpha_3 \sim 1 \times 10^{-4}, \quad \alpha_5 \sim 10^{-4} - 10^{-2}, \quad \alpha_6 \geq 0.1, \quad \alpha_9 \sim 3 \times 10^{-5}, \\ \alpha'_9 \sim 7 \times 10^{-6}, \quad \alpha_{10} \sim 8 \times 10^{-4}, \quad \alpha'_{10} \sim 2 \times 10^{-4}, \quad \alpha_{14} \sim 0.02. \end{aligned} \quad (74)$$

C Scalar potential and vacuum alignments

We present the scalar potential to discuss the vacuum alignment. The $SU(5) \times S_4 \times Z_4$ invariant superpotential is given as

$$\begin{aligned}
w = & \mu_1(\chi_1)_{1_1}^2 + \mu_2(\chi_2, \chi_3)_2^2 + \mu_3(\chi_4, \chi_5)_2^2 \\
& + \mu_4(\chi_9, \chi_{10}, \chi_{11})_{3_1}^2 + \mu_5(\chi'_9, \chi'_{10}, \chi'_{11})_{3_1}^2 + \mu_6(\chi_6, \chi_7, \chi_8)_{3_1} \otimes (\chi_{12}, \chi_{13}, \chi_{14})_{3_1} \\
& + \eta_1(\chi_1)_{1_1} \otimes (\chi_2, \chi_3)_2 \otimes (\chi_4, \chi_5)_2 + \eta_2(\chi_1)_{1_1} \otimes (\chi_6, \chi_7, \chi_8)_{3_1}^2 \\
& + \eta_3(\chi_1)_{1_1} \otimes (\chi_{12}, \chi_{13}, \chi_{14})_{3_1}^2 + \eta_4(\chi_1)_{1_1} \otimes (\chi_9, \chi_{10}, \chi_{11})_{3_1} \otimes (\chi'_9, \chi'_{10}, \chi'_{11})_{3_1} \\
& + \eta_5(\chi_2, \chi_3)_2^2 \otimes (\chi_4, \chi_5)_2 + \eta_6(\chi_2, \chi_3)_2 \otimes (\chi_6, \chi_7, \chi_8)_{3_1}^2 \\
& + \eta_7(\chi_2, \chi_3)_2 \otimes (\chi_9, \chi_{10}, \chi_{11})_{3_1} \otimes (\chi'_9, \chi'_{10}, \chi'_{11})_{3_1} + \eta_8(\chi_2, \chi_3)_2 \otimes (\chi_{12}, \chi_{13}, \chi_{14})_{3_1}^2 \\
& + \eta_9(\chi'_2, \chi'_3) \otimes (\chi_6, \chi_7, \chi_8) \otimes (\chi'_9, \chi'_{10}, \chi'_{11}) + \eta_{10}(\chi'_2, \chi'_3) \otimes (\chi_9, \chi_{10}, \chi_{11}) \otimes (\chi_{12}, \chi_{13}, \chi_{14}) \\
& + \eta_{11}(\chi_4, \chi_5)_2^3 + \eta_{12}(\chi_4, \chi_5)_2 \otimes (\chi_9, \chi_{10}, \chi_{11})_{3_1}^2 + \eta_{13}(\chi_4, \chi_5)_2 \otimes (\chi'_9, \chi'_{10}, \chi'_{11})_{3_1}^2 \\
& + \eta_{14}(\chi_4, \chi_5)_2 \otimes (\chi_6, \chi_7, \chi_8)_{3_1} \otimes (\chi_{12}, \chi_{13}, \chi_{14})_{3_1} + \eta_{15}(\chi_6, \chi_7, \chi_8)_{3_1}^2 \otimes (\chi'_9, \chi'_{10}, \chi'_{11})_{3_1} \\
& + \eta_{16}(\chi_9, \chi_{10}, \chi_{11})_{3_1}^3 + \eta_{17}(\chi_6, \chi_7, \chi_8)_{3_1} \otimes (\chi_9, \chi_{10}, \chi_{11})_{3_1} \otimes (\chi_{12}, \chi_{13}, \chi_{14})_{3_1} \\
& + \eta_{18}(\chi_9, \chi_{10}, \chi_{11})_{3_1} \otimes (\chi'_9, \chi'_{10}, \chi'_{11})_{3_1}^2 + \eta_{19}(\chi'_9, \chi'_{10}, \chi'_{11})_{3_1} \otimes (\chi_{12}, \chi_{13}, \chi_{14})_{3_1}^2, \tag{75}
\end{aligned}$$

which is written as

$$\begin{aligned}
w = & \mu_1\chi_1^2 + \mu_2(\chi_2^2 + \chi_3^2) + \mu_3(\chi_4^2 + \chi_5^2) + \mu_4(\chi_9^2 + \chi_{10}^2 + \chi_{11}^2) \\
& + \mu_5(\chi'_9{}^2 + \chi'_{10}{}^2 + \chi'_{11}{}^2) + \mu_6(\chi_6\chi_{12} + \chi_7\chi_{13} + \chi_8\chi_{14}) \\
& + \eta_1\chi_1(\chi_2\chi_4 + \chi_3\chi_5) + \eta_2\chi_1(\chi_6^2 + \chi_7^2 + \chi_8^2) + \eta_3\chi_1(\chi_{12}^2 + \chi_{13}^2 + \chi_{14}^2) \\
& + \eta_4\chi_1(\chi_9\chi'_9 + \chi_{10}\chi'_{10} + \chi_{11}\chi'_{11}) + \eta_5\{2\chi_2\chi_3\chi_4 + (\chi_2^2 - \chi_3^2)\chi_5\} \\
& + \eta_6\{\frac{\chi_2}{\sqrt{2}}(\chi_7^2 - \chi_8^2) + \frac{\chi_3}{\sqrt{6}}(-2\chi_6^2 + \chi_7^2 + \chi_8^2)\} \\
& + \eta_7\{\frac{\chi_2}{\sqrt{2}}(\chi_{10}\chi'_{10} - \chi_{11}\chi'_{11}) + \frac{\chi_3}{\sqrt{6}}(-2\chi_9\chi'_9 + \chi_{10}\chi'_{10} + \chi_{11}\chi'_{11})\} \\
& + \eta_8\{\frac{\chi_2}{\sqrt{2}}(\chi_{13}^2 - \chi_{14}^2) + \frac{\chi_3}{\sqrt{6}}(-2\chi_{12}^2 + \chi_{13}^2 + \chi_{14}^2)\} \\
& + \eta_9\{\frac{\chi'_2}{\sqrt{2}}(\chi_7\chi'_{10} - \chi_8\chi'_{11}) + \frac{\chi'_3}{\sqrt{6}}(-2\chi_6\chi'_9 + \chi_7\chi'_{10} + \chi_8\chi'_{11})\} \\
& + \eta_{10}\{\frac{\chi'_2}{\sqrt{2}}(\chi_{10}\chi_{13} - \chi_{11}\chi_{14}) + \frac{\chi'_3}{\sqrt{6}}(-2\chi_9\chi_{12} + \chi_{10}\chi_{13} + \chi_{11}\chi_{14})\} \\
& + \eta_{11}(3\chi_4^2\chi_5 - \chi_5^3) + \eta_{12}\{\frac{\chi_4}{\sqrt{2}}(\chi_{10}^2 - \chi_{11}^2) + \frac{\chi_5}{\sqrt{6}}(-2\chi_9^2 + \chi_{10}^2 + \chi_{11}^2)\} \\
& + \eta_{13}\{\frac{\chi_4}{\sqrt{2}}(\chi_{10}'^2 - \chi_{11}'^2) + \frac{\chi_5}{\sqrt{6}}(-2\chi_9'^2 + \chi_{10}'^2 + \chi_{11}'^2)\} \\
& + \eta_{14}\{\frac{\chi_4}{\sqrt{2}}(\chi_7\chi_{13} - \chi_8\chi_{14}) + \frac{\chi_5}{\sqrt{6}}(-2\chi_6\chi_{12} + \chi_7\chi_{13} + \chi_8\chi_{14})\} \\
& + \eta_{15}(\chi_7\chi_8\chi'_9 + \chi_6\chi_8\chi'_{10} + \chi_6\chi_7\chi'_{11}) + \eta_{16}\chi_9\chi_{10}\chi_{11} \\
& + \eta_{17}\{(\chi_7\chi_{11} + \chi_8\chi_{10})\chi_{12} + (\chi_6\chi_{11} + \chi_8\chi_9)\chi_{13} + (\chi_6\chi_{10} + \chi_7\chi_9)\chi_{14}\} \\
& + \eta_{18}(\chi_9\chi'_{10}\chi'_{11} + \chi_{10}\chi'_9\chi'_{11} + \chi_{11}\chi'_9\chi'_{10}) + \eta_{19}(\chi'_9\chi_{13}\chi_{14} + \chi'_{10}\chi_{12}\chi_{14} + \chi'_{11}\chi_{12}\chi_{13}). \tag{76}
\end{aligned}$$

VEVs of χ_i must be much larger than the weak scale. We assume that their VEVs are determined with neglecting supersymmetry breaking terms, i.e. $V_{\min} = 0$. The α_i and α'_j are very small except for α_6 , α_7 , α_8 , α_{14} , α'_2 and α'_3 . Then, conditions of the potential minimum, $V_{\min} = 0$ are written as

$$\begin{aligned}
\eta_2(\chi_6^2 + \chi_7^2 + \chi_8^2) + \eta_3\chi_{14}^2 &= 0, \\
\eta_6(\chi_7^2 - \chi_8^2) - \eta_8\chi_{14}^2 &= 0, \\
\frac{1}{\sqrt{6}}\eta_6(-2\chi_6^2 + \chi_7^2 + \chi_8^2) + \frac{1}{\sqrt{6}}\eta_8\chi_{14}^2 &= 0, \\
\frac{1}{\sqrt{6}}\eta_{14}\chi_8\chi_{14} &= 0, \\
\mu_6\chi_{14} &= 0, \\
\eta_{14}\chi_8\chi_{14} &= 0, \\
\eta_{17}\chi_7\chi_{14} &= 0, \\
\eta_{17}\chi_6\chi_{14} &= 0, \\
\eta_9\left(-\frac{1}{\sqrt{2}}\chi'_2 + \frac{1}{\sqrt{6}}\chi'_3\right) + \eta_{15}\chi_6\chi_7 &= 0, \\
\eta_{15}\chi_6\chi_8 &= 0, \\
-\frac{2}{\sqrt{6}}\eta_9\chi'_3\chi_6 + \eta_{15}\chi_7\chi_8 &= 0, \\
\eta_{10}\left(-\frac{1}{\sqrt{2}}\chi'_2 + \frac{1}{\sqrt{6}}\chi'_3\right)\chi_{14} &= 0, \\
\mu_6\chi_6 &= 0, \\
\mu_6\chi_7 &= 0,
\end{aligned} \tag{77}$$

where χ_i s are denoted as VEVs. Let us consider vacuum alignments of (χ_6, χ_7, χ_8) and (χ'_2, χ'_3) . Since we have a solution

$$\begin{aligned}
\eta_2(\chi_6^2 + \chi_7^2 + \chi_8^2) + \eta_3\chi_{14}^2 &= 0, \quad \eta_6(\chi_7^2 - \chi_8^2) = 0, \quad \frac{1}{\sqrt{6}}\eta_6(-2\chi_6^2 + \chi_7^2 + \chi_8^2) = 0, \\
\eta_8 &= \eta_9 = \eta_{10} = \eta_{14} = \eta_{15} = \eta_{17} = \mu_6 = 0,
\end{aligned} \tag{78}$$

the vacuum alignment $\chi_6 = \chi_7 = \chi_8$ is a possible solution. On the other hand, $\chi'_2 = \chi'_3$ is not guaranteed in Eq.(77). We may need another mechanism to realize the vacuum alignment of (χ'_2, χ'_3) [57].

References

- [1] T. Schwetz, M. Tortola, and J.W.F. Valle, arXiv:0808.2016;
G.L. Fogli, E. Lisi, A. Marrone, A. Palazzo, and A.M. Rotunno, Phys. Rev. Lett. **101** 141801 (2008); arXiv:0806.2649.
- [2] P.F. Harrison, D.H. Perkins, and W.G. Scott, Phys. Lett. B **530**, 167 (2002);
P.F. Harrison and W.G. Scott, Phys. Lett. B **535**, 163 (2002).
- [3] M. Fukugita, M. Tanimoto, and T. Yanagida, Phys. Rev. D **57**, 4429 (1998).
- [4] M. Tanimoto, Phys. Rev. D **59**, 017304 (1999), Phys. Lett. B **483**, 417 (2000).
- [5] P. F. Harrison and W. G. Scott, Phys. Lett. B **557**, 76 (2003), arXiv:hep-ph/0302025.
- [6] J. Kubo, A. Mondragon, M. Mondragon, and E. Rodriguez-Jauregui Prog. Theor. Phys. **109**, 795 (2003), Erratum-ibid. **114**, 287 (2005), arXiv: hep-ph/0302196.
- [7] S.-L. Chen, M. Frigerio, and E. Ma, Phys. Rev. D **70**, 073008 (2004),
arXiv:hep-ph/0404084.
- [8] W. Grimus and L. Lavoura, JHEP **08**, 013 (2005), arXiv:hep-ph/0504153.
- [9] F. Caravaglios and S. Morisi, arXiv:hep-ph/0503234.
- [10] R. N. Mohapatra, S. Nasri, and H.-B. Yu, Phys. Lett. B **639**, 318 (2006),
arXiv:hep-ph/0605020.
- [11] S. Morisi and M. Picariello, Int. J. Theor. Phys. **45**, 1267 (2006), arXiv:hep-ph/0505113.
- [12] N. Haba, A. Watanabe, and K. Yoshioka, Phys. Rev. Lett. **97** 041601 (2006),
arXiv:hep-ph/0603116.
- [13] S. Kaneko, H. Sawanaka, T. Shingai, M. Tanimoto, and K. Yoshioka, Prog. Theor. Phys. **117**, 161 (2007), arXiv:hep-ph/0609220.
- [14] Y. Koide, Eur. Phys. J. C **50**, 809 (2007), arXiv:hep-ph/0612058.
- [15] C.-Y. Chen and L. Wolfenstein, Phys. Rev. D **77**, 093009 (2008), arXiv:0709.3767.
- [16] F. Feruglio and Y. Lin, Nucl. Phys. B **800**, 77 (2008), arXiv:0712.1528.
- [17] A. Mondragon, M. Mondragon, and E. Peinado, AIP Conf. Proc. **1026**, 164 (2008),
arXiv:0712.2488.
- [18] M. Mitra and S. Choubey, arXiv:0806.3254.
- [19] W. Grimus and L. Lavoura, arXiv:0811.4766.
- [20] E. Ma and G. Rajasekaran, Phys. Rev. D **64**, 113012 (2001), arXiv:hep-ph/0106291;
E. Ma, Mod. Phys. Lett. A **17**, 2361 (2002).

- [21] K. S. Babu, E. Ma, and J. W. F. Valle, Phys. Lett. B **552**, 207 (2003), arXiv:hep-ph/0206292.
- [22] M. Hirsch, J.C. Romao, S. Skadhauge, J.W.F. Valle, and A. Villanova del Moral, Phys. Rev. D **69**, 093006 (2004), arXiv:hep-ph/0312265.
- [23] E. Ma, Phys. Rev. D **70**, 031901 (2004), arXiv:hep-ph/0404199.
- [24] G. Altarelli and F. Feruglio, Nucl. Phys. B **720**, 64 (2005), arXiv:hep-ph/0504165; Nucl. Phys. B **741**, 215 (2006), arXiv:hep-ph/0512103.
- [25] S.-L. Chen, M. Frigerio, and E. Ma, Nucl. Phys. B **724**, 423 (2005), arXiv:hep-ph/0504181.
- [26] A. Zee, Phys. Lett. B **630**, 58 (2005), arXiv:hep-ph/0508278.
- [27] B. Adhikary, B. Brahmachari, A. Ghosal, E. Ma, and M. K. Parida, Phys. Lett. B **638**, 345 (2006), arXiv:hep-ph/0603059.
- [28] J. W. F. Valle, J. Phys. Conf. Ser. **53**, 473 (2006), arXiv:hep-ph/0608101.
- [29] X.-G. He, Y.-Y. Keum, and R.R. Volkas, JHEP **0604**, 039 (2006).
- [30] E. Ma, H. Sawanaka, and M. Tanimoto, Phys. Lett. B **641**, 301 (2006).
- [31] B. Adhikary and A. Ghosal, Phys. Rev. D **75**, 073020 (2007), arXiv:hep-ph/0609193.
- [32] E. Ma, Phys. Rev. D **70**, 031901 (2004), Phys. Rev. D **72**, 037301 (2005), Mod. Phys. Lett. A **22**, 101 (2007), arXiv:hep-ph/0610342.
- [33] G. Altarelli, F. Feruglio, and Y. Lin, Nucl. Phys. B **775**, 31 (2007), arXiv:hep-ph/0610165.
- [34] S. F. King and M. Malinský, Phys. Lett. B **645**, 351 (2007).
- [35] M. Hirsch, A. S. Joshipura, S. Kaneko, and J.W.F. Valle, Phys. Rev. Lett. **99**, 151802 (2007), arXiv:hep-ph/0703046.
- [36] L. Lavoura and H. Kühböck, Mod. Phys. Lett. A **22**, 181 (2007), arXiv:0711.0670.
- [37] M. Honda and M. Tanimoto, Prog. Theor. Phys. **119**, 585 (2008), arXiv:0801.0181.
- [38] F. Bazzocchi, S. Kaneko, and S. Morisi, JHEP **03**, 063 (2008), arXiv:0707.3032.
- [39] F. Bazzocchi, M. Frigerio, and S. Morisi, arXiv:0809.3573.
- [40] Y. Lin, arXiv:0804.2867.
- [41] M. Hirsch, S. Morisi, and J.W.F. Valle, arXiv:0810.0121.
- [42] F. Feruglio, C. Hagedorn, Y. Lin, and L. Merlo, Nucl. Phys. B **775**, 120 (2007), arXiv:hep-ph/0702194.

- [43] M. C. Chen and K. T. Mahanthappa, Phys. Lett. B **652**, 34 (2007), arXiv:0705.0714.
- [44] A. Aranda, Phys. Rev. D **76**, 111301 (2007), arXiv:0707.3661.
- [45] D. A. Eby, P. H. Frampton, and S. Matsuzaki, arXiv:0810.4899.
- [46] P. H. Frampton and T. W. Kephart, Int. J. Mod. Phys. A **10**, 4689 (1995), arXiv:hep-ph/9409330.
- [47] G. J. Ding, Phys. Rev. D **78**, 036011 (2008).
- [48] C. S. Lam, arXiv:0809.1185.
- [49] F. Bazzocchi and S. Morisi, arXiv:0811.0345.
- [50] E. Ma, Phys. Lett. B **632**, 352 (2006), arXiv:hep-ph/0508231.
- [51] C. Hagedorn, M. Lindner, and R. N. Mohapatra, JHEP **06**, 042 (2006), arXiv:hep-ph/0602244.
- [52] Y. Cai and H.-B. Yu, Phys. Rev. D **74**, 115005 (2006), arXiv:hep-ph/0608022.
- [53] H. Zhang, Phys. Lett. B **655**, 132 (2007), arXiv:hep-ph/0612214.
- [54] Y. Koide, JHEP **08**, 086 (2007), arXiv:0705.2275.
- [55] H. Fritzsch, Phys. Lett. B **73**, 317 (1978); Nucl. Phys. B **115**, 189 (1979).
- [56] H. Fusaoka and Y. Koide, Phys. Rev. D **57**, 3986 (1998), hep-ph/9712201.
- [57] T. Kobayashi, Y. Omura, and K. Yoshioka, arXiv:0809.3064; G. Seidl, arXiv:0811.3775.